


SN 09/980,869
TSRI 696.1

and R^3 is $-\text{NHC(O)}R^4$, wherein R^4 is a hydrocarbon having between 1 and 20 carbon atoms.

33. (once amended) An inhibitor of hexoaminidase or glycosidase represented by the following structure:



wherein R^1 is methyl sulfate; R^2 is selected from the group consisting of hydrogen, methyl, ethyl, and a branched or unbranched hydrocarbon of between 3 and 8 carbon atoms; and R^3 is $-\text{NHC(O)}R^4$, wherein R^4 is a hydrocarbon having between 1 and 20 carbon atoms.

REMARKS

The above amendments are submitted by the Applicant in order to correct an error of indefiniteness.

Respectfully submitted,

Donald G. Lewis
Reg. No. 28,636
The Scripps Research Institute
10550 N. Torrey Pines Road TPC-8
San Diego, CA 92037
July 23, 2003
(858) 784-2937

- 2 -

TSRI 696.1
SN 09/980,869


APPENDIX

VERSION OF CLAIMS
WITH MARKINGS TO SHOW CHANGES MADE

A marked-up version of amendments to Claims 30 and 33 is provided:


In the Claims:

- 5 30. (once amended) An inhibitor of hexoaminidase or glycosidase represented by the following structure:



10 wherein R^1 is sulfate; R^2 is selected from the group consisting of hydrogen, methyl, ethyl, and a branched or unbranched hydrocarbon of between 3 and 8 carbon atoms; and R^3 is $-NHC(O)R^4$, wherein R^4 is a hydrocarbon having between 1 and 20 carbon atoms.

- 15 33. (once amended) An inhibitor of hexoaminidase or glycosidase represented by the following structure:



20 wherein R^1 is methyl sulfate; R^2 is selected from the group consisting of hydrogen, methyl, ethyl, and a branched or unbranched hydrocarbon of between 3 and 8 carbon atoms; and R^3 is $-NHC(O)R^4$, wherein R^4 is a hydrocarbon having between 1 and 20 carbon atoms.